

Texas A&M University-San Antonio

Digital Commons @ Texas A&M University-San Antonio

Mathematics Faculty Publications

College of Arts and Sciences

12-2022

Optimal Test Plan of Step Stress Partially Accelerated Life Testing for Alpha Power Inverse Weibull Distribution under Adaptive Progressive Hybrid Censored Data and Different Loss Functions

Refah Alotaibi

Ehab M. Almetwally

Qiuchen Hai

Hoda Rezk




Follow this and additional works at: https://digitalcommons.tamusa.edu/math_faculty



Part of the [Mathematics Commons](#)

Article

Optimal Test Plan of Step Stress Partially Accelerated Life Testing for Alpha Power Inverse Weibull Distribution under Adaptive Progressive Hybrid Censored Data and Different Loss Functions

Refah Alotaibi ¹, Ehab M. Almetwally ^{2,3}, Qiuchen Hai ⁴ and Hoda Rezk ^{5,*}

¹ Department of Mathematical Sciences, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia

² Department of Mathematical Statistical, Faculty of Graduate Studies for Statistical Research, Cairo University, Cairo 12613, Egypt

³ Department of Statistical, Faculty of Business Administration, Delta University for Science and Technology, Talkha 7730103, Egypt

⁴ Department of Mathematical, Physical, and Engineering Sciences, Texas A&M University-San Antonio, San Antonio, TX 78224, USA

⁵ Department of Statistics, Al-Azhar University, Cairo 4434003, Egypt

* Correspondence: hodaragab2009@yahoo.com



Citation: Alotaibi, R.; Almetwally, E.M.; Hai, Q.; Rezk, H. Optimal Test Plan of Step Stress Partially Accelerated Life Testing for Alpha Power Inverse Weibull Distribution under Adaptive Progressive Hybrid Censored Data and Different Loss Functions. *Mathematics* **2022**, *10*, 4652. <https://doi.org/10.3390/math10244652>

Academic Editors: Alexandru Agapie, Denis Enachescu, Vlad Stefan Barbu and Bogdan Iftimie

Received: 4 October 2022

Accepted: 3 December 2022

Published: 8 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Accelerated life tests are used to explore the lifetime of extremely reliable items by subjecting them to elevated stress levels from stressors to cause early failures, such as temperature, voltage, pressure, and so on. The alpha power inverse Weibull (APIW) distribution is of great significance and practical applications due to its appealing characteristics, such as its flexibilities in the probability density function and the hazard rate function. We analyze the step stress partially accelerated life testing model with samples from the APIW distribution under adaptive type II progressively hybrid censoring. We first obtain the maximum likelihood estimates and two types of approximate confidence intervals of the distributional parameters and then derive Bayes estimates of the unknown parameters under different loss functions. Furthermore, we analyze three probable optimum test techniques for identifying the best censoring under different optimality criteria methods. We conduct simulation studies to assess the finite sample performance of the proposed methodology. Finally, we provide a real data example to further demonstrate the proposed technique.

Keywords: the alpha power inverse Weibull distribution; step stress partially accelerated life testing; adaptive progressive hybrid censored data; loss functions

MSC: 65C20; 60E05; 62P30; 62L15

1. Introduction

The reliability of products has recently grown greatly in the present era of technical achievements due to an ongoing effort for improving manufacturing processes in various companies. Under the presence of high competition to launch their products within a short time period, direct use of traditional life testing methodologies will be an expensive and time-consuming operation for evaluating the lifetime of a product to predict product failures. As a result, accelerated life tests (ALTs) are usually employed to explore the lifetime of extremely reliable products, as they can be used with elevated stress levels of stressors to trigger early failures, such as temperature, voltage (electric field), pressure, and so on. Thereafter, the constant-stress and step-stress models in the ALTs have been studied in life testing and reliability analyses; see, for example, [1–4].

It is known that each product sample in the ALTs is typically analyzed under a constant-stress scenario, subjected to some continuous amounts of constant stress until

all units fail or the test is cancelled for any reason, such as censoring plan. However, the test conditions associated with step-stress models do not remain constant throughout the tests, since the stress on a sample of test units could increase step by step at a prescribed period or concurrently when a fixed number of failures occurs. In addition, the ALTs often use a suitable physical model to extrapolate the collected breakdown information under accelerated settings, whereas it is difficult to select a proper physical model to describe the life stress relationships in practical situations [5]. To overcome these drawbacks of ALTs, researchers may employ partially accelerated life testing (PALT), which is classified into two types: constant-stress loading and step-stress loading. In the constant-stress PALT (CSPALT), each sample of tested items is subjected to normal and accelerated levels of constant stress until all units fail or the test is terminated; see, for example, [6,7]. In the step-stress PALT (SSPALT), certain objects or materials are initially tested under normal or usage settings for a predetermined amount of time before being subjected to accelerated test conditions until the termination time; see, for example, [8–10].

In life-testing and reliability trials, data are commonly censored due to time and cost constraints. The hybrid censoring scheme [11], which includes Type-I and Type-II censorings as special cases, is commonly utilized in reliability analysis. We, here, refer the interested reader to [12] for a nice overview of the hybrid censoring. However, the hybrid censoring scheme lacks an option to delete units during the testing period due to time and cost constraints. To address this issue, a progressive censoring scheme was developed by allowing for the deletion of experimental units at various periods of time throughout the test; see, for example, [13,14] in detail. It is worth pointing out that in the progressively Type-II hybrid censoring, the number of required failures and the number of items that must be deleted are determined in advance, whereas there is no time constraint on the experiment, leading to a very long period.

To address this issue, [15] proposed the Type-I Progressive Hybrid Censoring Scheme (TIPHCS), with an additional time and failure constraint that the experiment will run until a predetermined time point or a predetermined number of failures, whichever comes first. However, since the sample size in TIPHCS is random, only a few or even no failure would occur before a pre-specified time limit, resulting in poor efficiency of the parameter estimation. The authors of [16] proposed an adaptive type-II PHCS (AT-II PHCS), in which n units are placed on a life test with a predetermined number of failures m and a pre-fixed progressive censoring scheme $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m$, but the experimenter is allowed to change some of the ε_{i_w} s during the experiment depending on situations. At the initial failure time, $z_{1:m:n}$, ε_1 units are randomly selected from the remaining $n - 1$ alive items and are then removed from the experiment. At the second failure time, $z_{2:m:n}$, ε_2 units of the remaining $n - 1 - \varepsilon_1$ units are eliminated at random, and so on. If the m -th failure time $z_{m:m:n}$, occurs before the predetermined time δ , all the remaining $\varepsilon_m = n - m - \sum_{i=1}^m \varepsilon_i$ units are removed and the experiment terminates at time $z_{m:m:n}$. The AT-II PHCS allows the experiment to run over the test termination time restriction. As a result, if $z_{m:m:n} > \delta$, the experiment will soon be stopped by setting $\varepsilon_{c+1}, \varepsilon_{c+2}, \dots, \varepsilon_{m-1} = 0$. This means that if $z_{c:m:n} < \delta < z_{c+1:m:n}$, with $c + 1 < m$ and $y_{c:m:n}$ is the c -th failure time that occurred before δ , no surviving item will be removed from the experiment until the effective sample of m failures is attained, resulting in the remaining units $\varepsilon_m = n - c - \sum_{i=1}^c \varepsilon_i$.

Due to the importance of the AT-II PHCS, numerous authors have investigated the problem of parameter estimation in different statistical models based on this censoring scheme; see, for example, Refs. [17–19] for the Weibull distribution, Ref. [20] for the log-normal distribution, Ref. [21] for the exponentiated Weibull distribution, Refs. [22,23] for the extended Weibull distribution, Ref. [24] for the Burr Type-XII distribution, Ref. [25] for the exponentiated Pareto distribution, Ref. [26] for the inverted NH distribution, Ref. [27] for the Weibull generalized exponential distribution, Ref. [28] for the exponentiated exponential distribution, Ref. [29] for the exponentiated power Lindley distribution, and references cited therein. To the best of our knowledge, little research attention has been devoted to the

alpha power inverse Weibull (APIW) distribution [30]. This observation motivates us to investigate statistical inference of the APIW distribution under AT-II PHCS.

Due to flexibilities in its probability density function (PDF) and hazard rate function (HRF), the APIW distribution has become a useful model in the study of life testing and reliability analyses. The cumulative distribution function (CDF), PDF, survival function (SF), and HRF for an APIW random variable T are given by

$$F(t; \alpha, \beta, \theta) = \frac{\alpha e^{-\beta t^{-\theta}} - 1}{\alpha - 1}, \quad \alpha, \beta, \theta, t > 0, \tag{1}$$

$$f(t; \alpha, \beta, \theta) = \frac{\log(\alpha) \theta \beta e^{-\beta t^{-\theta}} t^{-\theta-1} \alpha e^{-\beta t^{-\theta}}}{\alpha - 1}, \quad \alpha, \beta, \theta, t > 0, \tag{2}$$

$$S(t; \alpha, \beta, \theta) = \frac{\alpha}{\alpha - 1} \left(1 - \alpha e^{-\beta t^{-\theta}} - 1 \right), \quad \alpha, \beta, \theta, t > 0, \tag{3}$$

and

$$h(t; \alpha, \beta, \theta) = \frac{\log(\alpha) \theta \beta e^{-\beta t^{-\theta}} t^{-\theta-1} \alpha e^{-\beta t^{-\theta}} - 1}{\left(1 - \alpha e^{-\beta t^{-\theta}} - 1 \right)}, \quad \alpha, \beta, \theta, t > 0, \tag{4}$$

respectively, where $\alpha > 0$ and $\theta > 0$ are the shape parameters and $\beta > 0$ is a scale parameter. This distribution includes many well-known distributions as special cases, such as the alpha power Fréchet, alpha power inverse Rayleigh, alpha power inverse exponential, inverse Weibull, Fréchet, inverse Rayleigh, and the inverse exponential distributions. In addition, it has closed-form expressions of the SF and HRF, which make the distribution a good alternative to commonly used distributions in life-testing analysis.

In this paper, we analyze the step stress partially accelerated life testing model with samples from the APIW distribution under the AT-II PHCS. We first consider the MLEs and derive asymptotic confidence interval and bootstrap confidence intervals of the model parameters. We then propose Bayes estimates of the unknown parameters with non-informative and informative priors under the symmetric and asymmetric loss functions. In addition, we identify the best progressive censoring scheme to the most information about the unknown parameters among all conceivable progressive censoring schemes. Numerical results from simulation studies and a real-data application show that the performance of the proposed technique is quite satisfactory for analyzing censored data under different sampling schemes.

The rest of this paper is organized as follows. Section 2 describes the lifetime model and the test assumptions. Section 3 derives the MLEs of the APIW parameters under the AT-II PHCS. Section 4 constructs the confidence intervals of the unknown parameters. Bayesian analysis of the unknown parameters is provided in Section 5. We carry out simulations in Section 6 to investigate the finite sample performance of the proposed model. In Section 7, a real-data example is provided for illustrative purposes. Finally, concluding remarks are provided in Section 8 with Fisher information of the model deferred to the Appendix A.

2. Assumptions and Procedure for Testing

Suppose that in a simple SSPALT, the test employs only two stress levels, S_u (normal operating circumstances) and S_a (accelerated condition), such that $S_u < S_a$, where S_u and S_a are twins. Under each stress level, at least one failure should occur. We assume that at both stress levels, the failures of the test items follow the APIW distribution in (2). Then, the lifetime Z of a test item follows a TRV model given by

$$Z = \begin{cases} T, & \text{if } T < \tau \\ \tau + \frac{T-\tau}{\lambda}, & T > \tau \end{cases}$$

where T indicates the lifetime of an item under the stress ST and represents the time point at which stress ST is switched from u to a , and $\lambda > 1$ is an accelerated factor (AF). Then, under the TRV model, we obtain the PDF, CDF, and SF of Z given by

$$f_u(z; \alpha, \beta, \theta) = \frac{\log(\alpha) \theta \beta e^{-\beta z^{-\theta}} z^{-\theta-1} \alpha^{e^{-\beta z^{-\theta}}}}{\alpha - 1}, \tag{5}$$

$$F_u(z; \alpha, \beta, \theta) = \frac{\alpha^{e^{-\beta z^{-\theta}}} - 1}{\alpha - 1}, \tag{6}$$

$$S_u(z; \alpha, \beta, \theta) = \frac{\alpha}{\alpha - 1} \left(1 - \alpha^{e^{-\beta z^{-\theta}}} - 1 \right) \tag{7}$$

Now, at stress $ST = a$, the PDF, CDF, and SF of Z are produced as follows:

$$f_a(z; \alpha, \beta, \theta, \lambda) = \frac{\log(\alpha) \theta \beta e^{-\beta[\tau + \lambda(z-\tau)]^{-\theta}} [\tau + \lambda(z-\tau)]^{-\theta-1} \alpha^{e^{-\beta[\tau + \lambda(z-\tau)]^{-\theta}}}}{\alpha - 1}, \tag{8}$$

$$F_a(z; \alpha, \beta, \theta, \lambda) = \frac{\alpha^{e^{-\beta[\tau + \lambda(z-\tau)]^{-\theta}}} - 1}{\alpha - 1}, \alpha, \beta, \theta, z > 0, \lambda > 1, \tag{9}$$

$$S_a(z; \alpha, \beta, \theta, \lambda) = \frac{\alpha}{\alpha - 1} \left(1 - \alpha^{e^{-\beta[\tau + \lambda(z-\tau)]^{-\theta}}} - 1 \right), \tag{10}$$

where $\alpha, \beta, \theta, z > 0, \lambda > 1$. We assume that a sample of n items is assigned to the stress level S_u to test according to the SSPALT and a known progressive censoring scheme $\epsilon_1, \epsilon_2, \dots, \epsilon_m$. The test will proceed and the items from n that do not fail up to time S_u are placed through S_a to test, and the test will continue until the censorship time is reached. If the $m - th$ failure does not occur within the censoring point δ , no item will be removed from the test. The testing will continue until the $m - th$ failure is registered, at which point it will be terminated when all remaining items are eliminated. As a result, the implemented scheme is $\epsilon_1, \epsilon_2, \dots, \epsilon_c, 0, 0, \dots, 0, \epsilon_m$. Thus, we obtain the observed samples given by

$$z_{1:m:n} < z_{2:m:n} < \dots < z_{m_u:m:n} < \tau < z_{m_u+1:m:n} < \dots < z_{c:m:n} < \delta < z_{c+1:m:n} < \dots < z_{m:m:n},$$

illustrated in Figure 1.




Figure 1. Illustration of the AT-II PHCS scheme.

We observe from this figure that the AT-II PHCS is a special instance of the AT-II PHCS as $\delta \rightarrow 0$ and that the AT-II PHCS reduces to the classical Type-II PHCS as $\delta \rightarrow \infty$.

3. The Parameter Estimation

The resulting likelihood function of the data under the AT-II PHCS is given by

$$L(z; \alpha, \beta, \theta, \lambda) \propto \prod_{i=1}^{m_u} \{f_u(z_i)[R_u(z_i)]^{\varepsilon_i}\} \prod_{i=m_u+1}^m \{f_a(z_i)[R_a(z_i)]^{\varepsilon_i}[R_a(z_m)]^{\varepsilon_m}\}, \tag{11}$$

where $z_i = z_{i:m:n}$, $\varepsilon_m = n - m - \sum_{i=1}^c \varepsilon_i$. Then it follows

$$L(z; \alpha, \beta, \theta, \lambda) \propto \prod_{i=1}^{m_u} \left\{ \frac{\log(\alpha) \theta \beta e^{-\beta z_i^{-\theta}} z_i^{-\theta-1} \alpha^{e^{-\beta z_i^{-\theta}}}}{\alpha - 1} \left[\frac{\alpha}{\alpha - 1} \left(1 - \alpha^{e^{-\beta z_i^{-\theta}} - 1} \right) \right]^{\varepsilon_i} \right\} \\ \times \prod_{i=m_u+1}^m \left\{ \frac{\log(\alpha) \theta \beta e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} [\tau+\lambda(z_i-\tau)]^{-\theta-1} \alpha^{e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}}}}{\alpha - 1} \left[\frac{\alpha}{\alpha - 1} \left(1 - \alpha^{e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1} \right) \right]^{\varepsilon_i} \right. \\ \left. \left[\frac{\alpha}{\alpha - 1} \left(1 - \alpha^{e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1} \right) \right]^{\varepsilon_m} \right\} \tag{12}$$

The MLE is commonly used to estimate the unknown parameters, as it effectively and efficiently yields estimates with good statistical properties. By taking the natural logarithm on both sides of Equation (12), we obtain the log-likelihood equation $L(z; \alpha, \beta, \theta, \lambda) = \ell$ as follows

$$\ell = m_u \log(\log(\alpha)) - m_u \log(\alpha - 1) + m_u \log(\theta) + m_u \log(\beta) - \beta \sum_{i=1}^{m_u} z_i^{-\theta} - (\theta + 1) \sum_{i=1}^{m_u} \log(z_i) \\ + \log(\alpha) \sum_{i=1}^{m_u} e^{-\beta z_i^{-\theta}} + \sum_{i=1}^{m_u} \varepsilon_i \log(\alpha) - \sum_{i=1}^{m_u} \varepsilon_i \log(\alpha - 1) + \sum_{i=1}^{m_u} \varepsilon_i \left(1 - \alpha^{e^{-\beta z_i^{-\theta}} - 1} \right) \\ + m \log(\log(\alpha)) - m \log(\alpha - 1) + m \log(\theta) + m \log(\beta) - \beta \sum_{i=m_u+1}^m [\tau + \lambda(z_i - \tau)]^{-\theta} \\ - (\theta + 1) \sum_{i=m_u+1}^m \log[\tau + \lambda(z_i - \tau)] + \log(\alpha) \sum_{i=m_u+1}^m e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} + \sum_{i=m_u+1}^m (\varepsilon_i) \log(\alpha) \\ - \sum_{i=m_u+1}^m (\varepsilon_i) \log(\alpha - 1) + \sum_{i=m_u+1}^m \varepsilon_i \log \left(1 - \alpha^{e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1} \right) + \varepsilon_m \log(\alpha) - \varepsilon_m \log(\alpha - 1) \\ + \varepsilon_m \log \left(1 - \alpha^{e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1} \right) \tag{13}$$

The MLEs of the parameters α, β, θ and λ can be obtained by solving the following nonlinear system equations

$$\frac{\partial \ell}{\partial \alpha} = \frac{m_u + m}{\alpha \log \alpha} - \frac{(m_u + m)}{(\alpha - 1)} + \frac{1}{\alpha} \sum_{i=1}^{m_u} e^{-\beta z_i^{-\theta}} - \frac{1}{\alpha} \sum_{i=1}^{m_u} \varepsilon_i - \frac{1}{\alpha - 1} \sum_{i=1}^{m_u} \varepsilon_i - \log(\alpha) \sum_{i=1}^{m_u} \varepsilon_i \alpha^{e^{-\beta z_i^{-\theta}} - 1} \\ + \frac{1}{\alpha} \sum_{i=m_u+1}^m \varepsilon_i e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} + \frac{1}{\alpha} \sum_{i=m_u+1}^m (\varepsilon_i) + \frac{1}{(\alpha - 1)} \sum_{i=m_u+1}^m (\varepsilon_i) \\ + \log(\alpha) \sum_{i=m_u+1}^m \varepsilon_i \frac{\alpha^{e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1}}{\left(1 - \alpha^{e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1} \right)} + \frac{\varepsilon_m}{\alpha} - \frac{\varepsilon_m}{(\alpha - 1)} + \frac{\varepsilon_m \log(\alpha) \alpha^{e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1}}{1 - \alpha^{e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1}}, \tag{14}$$

$$\begin{aligned}
 & \frac{\partial \ell}{\partial \theta} \\
 &= \frac{m_u+m}{\theta} + \beta \sum_{i=1}^{m_u} z_i^{-\theta} \log(z_i) - \sum_{i=1}^{m_u} \log(z_i) \\
 &+ \log(\alpha) \sum_{i=1}^{m_u} \beta z_i^{-\theta} \log(z_i) e^{-\beta z_i^{-\theta}} + \sum_{i=1}^{m_u} e^{-\beta z_i^{-\theta}} \beta z_i^{-\theta} \log(z_i) \left(e^{-\beta z_i^{-\theta}} - 1 \right) \varepsilon_i \alpha e^{-\beta z_i^{-\theta} - 2} \\
 &+ \beta \sum_{i=m_u+1}^m [\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] \\
 &- \sum_{i=m_u+1}^m \log[\tau + \lambda(z_i - \tau)] + \log(\alpha) \sum_{i=m_u+1}^m e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \beta [\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] \\
 &+ \sum_{i=m_u+1}^m \frac{\varepsilon_i e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \beta [\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta} - 2} \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)} \\
 &+ \frac{\varepsilon_m e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \beta [\tau + \lambda(z_m - \tau)]^{-\theta} \log[\tau + \lambda(z_m - \tau)] \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta} - 2} \right)}{\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1},
 \end{aligned} \tag{15}$$

$$\begin{aligned}
 \frac{\partial \ell}{\partial \beta} &= \frac{m_u}{\beta} - \sum_{i=1}^{m_u} z_i^{-\theta} - \log(\alpha) \sum_{i=1}^{m_u} z_i^{-\theta} e^{-\beta z_i^{-\theta}} - \sum_{i=1}^{m_u} \varepsilon_i z_i^{-\theta} e^{-\beta z_i^{-\theta}} \left(\alpha e^{-\beta z_i^{-\theta}} - 1 \right) + \frac{m}{\beta} + \sum_{i=m_u+1}^m [\tau + \lambda(z_i - \tau)]^{-\theta} \\
 &+ \log(\alpha) \sum_{i=m_u+1}^m [\tau + \lambda(z_i - \tau)]^{-\theta} e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \\
 &+ \sum_{i=m_u+1}^m \frac{\varepsilon_i \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)}{1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1} \\
 &+ \frac{\varepsilon_m \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) [\tau + \lambda(z_m - \tau)]^{-\theta} \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)}{1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1},
 \end{aligned} \tag{16}$$

$$\begin{aligned}
 & \frac{\partial \ell}{\partial \lambda} \\
 &= -\beta \sum_{i=m_u+1}^m (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta - 1} - \sum_{i=m_u+1}^m \frac{(\theta + 1)(z_i - \tau)}{[\tau + \lambda(z_i - \tau)]} \\
 &+ \beta \log(\alpha) \sum_{i=m_u+1}^m e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta - 1} \\
 &+ \frac{\sum_{i=m_u+1}^m \varepsilon_i e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \beta (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta - 1} \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta} - 2}}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)} \\
 &+ \frac{\varepsilon_m e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \beta (z_m - \tau) [\tau + \lambda(z_m - \tau)]^{-\theta - 1} \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta} - 2}}{1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1}.
 \end{aligned} \tag{17}$$

We observe that it is difficult to get closed-form solutions of the parameters from the above nonlinear equations. As a result, we employ an iterative approach, such as Newton–Raphson, to find numerical solutions to the nonlinear systems.

4. Confidence Intervals

A confidence interval (CI) is a collection of numbers that serves as reasonable approximations to an unknown population characteristic (e.g., [31]). We consider two types of CIs for the unknown parameters as follows.

4.1. Approximate Confidence Intervals

According to large sample theory, the MLEs are consistent and regularly distributed under certain regularity conditions. To be more specific, $[(\hat{\alpha} - \alpha), (\hat{\theta} - \theta), (\hat{\beta} - \beta), (\hat{\lambda} - \lambda)] \sim N(0, \sigma)$ is known to yield the asymptotic distribution of MLEs of α, θ, β and λ , where $\sigma = \sigma_{ij}, i, j = 1, 2, 3, 4$ is the unknown parameters’ in the variance–covariance matrix. The

inverse of the Fisher information matrix is an estimate of the variance–covariance matrix. The estimated $100(1 - \pi)\%$ two-sided CIs for the unknown parameter are provided by

$$(\hat{\pi}_{iL}, \hat{\pi}_{iU}) : \hat{\pi}_i \mp z_{1-\frac{\pi}{2}} \sqrt{\hat{\sigma}_{ij}}, i = 1, 2, 3, 4,$$

where $z_{1-\pi/2}$ is the π -th percentile of the standard normal distribution. However, the above asymptotic CIs may not perform well due to an asymmetric property of the APIW distribution. To deal with this issue, we consider the parametric bootstrap percentile intervals as an alternative [32].

4.2. Bootstrap Confidence Intervals

We consider the parametric bootstrap sampling with percentile intervals, which can be implemented using Algorithm 1.

Algorithm 1. Bootstrap

1. Step 0, basic setup:
 2. Set $b = 1$
 3. Determine the MLE values of $\omega = (\alpha, \theta, \beta, \lambda)$, as showing by $\hat{\omega} = (\hat{\alpha}, \hat{\theta}, \hat{\beta}, \hat{\lambda})$.
 4. Step 1: Sample
 5. Get the b th bootstrap resample t_p^* from $F(\cdot|\hat{\omega})$, where is the MLE from Step 0.
 6. Step 2: Estimates from the bootstrap:
 7. Calculate the b th bootstrap estimations.
 8. $\hat{\omega}^{*b} = (\hat{\alpha}^{*b}, \hat{\theta}^{*b}, \hat{\beta}^{*b}, \hat{\lambda}^{*b})$,
 9. Utilize the t_p^* resample obtained in Step 1.
 10. Step three, repeat:
 11. Set $b \leftarrow b+1$,
 12. Steps 1–3 are then repeated until $b = G$.
 13. Step 4: In ascending sequence, begin:
 14. Sort the estimates in increasing order so that
 15. $\{\hat{\alpha}^{*[1]}, \hat{\alpha}^{*[2]}, \dots, \hat{\alpha}^{*[G]}\}, \{\hat{\theta}^{*[1]}, \hat{\theta}^{*[2]}, \dots, \hat{\theta}^{*[G]}\}, \{\hat{\beta}^{*[1]}, \hat{\beta}^{*[2]}, \dots, \hat{\beta}^{*[G]}\}, \{\hat{\lambda}^{*[1]}, \hat{\lambda}^{*[2]}, \dots, \hat{\lambda}^{*[G]}\}$
-

The $100(1 - \omega)\%$ percentile bootstrap CIs for the unknown parameter are computed as follows

$$(\hat{\omega}_{iL}, \hat{\omega}_{iU}) = (\hat{\omega}_i^{*[\frac{\pi}{2}]G}, \hat{\omega}_i^{*[(1-\frac{\pi}{2})G]}), i = 1, 2, 3, 4,$$

where $\hat{\omega}_1^* = \alpha^*$, $\hat{\omega}_2^* = \theta^*$, $\hat{\omega}_3^* = \beta^*$, and $\hat{\omega}_4^* = \lambda^*$.

5. Bayesian Estimation

In this section, we focus on Bayes estimation for the unknown parameters. Bayesian analysis begins with prior specifications for the unknown parameters. In this paper, we assume that the parameters α , θ , β , and λ are statistically independent and follow independent gamma distributions, denoted by $gamma(a_j, b_j); j = 1, \dots, 4$, respectively. The joint priors for the APIW parameters can be written as

$$\varphi(\alpha, \theta, \beta, \lambda) \propto \alpha^{a_1-1} e^{-b_1\alpha} \theta^{a_2-1} e^{-b_2\theta} \beta^{a_3-1} e^{-b_3\beta} \lambda^{a_4} e^{-b_4\lambda}, \tag{18}$$

where $a_j \geq 0$ and $b_j \geq 0; j = 1, \dots, 4$ are pre-determined hyperparameters that reflect prior knowledge of the unknown parameters. The resulting joint posterior distribution of the unknown parameters is given by

$$L(\alpha, \theta, \beta, \lambda | \underline{t}) \propto \varphi(\alpha, \theta, \beta, \lambda) \prod_{i=1}^4 \prod_{j=1}^{n_i} f(t_{ij}) (1 - F(t_{ij}))^{d_i}, \tag{19}$$

which is usually unidentifiable. Thus, we employ Markov chain Monte Carlo (MCMC) methods to generate posterior samples of the unknown parameters for making posterior inference. In particular, the acquired samples will also be used to approximate Bayes estimates and obtain the corresponding highest posterior density (HPD) credible ranges for

the unknown parameters [33]. In this paper, we obtain the Bayes estimates of the unknown parameters under the symmetric (SLF) and asymmetric (ELF) loss functions, which are denoted as

$$\ell(\alpha, \tilde{\alpha}) = (\tilde{\alpha} - \alpha)^2, \ell(\theta, \tilde{\theta}) = (\tilde{\theta} - \theta)^2, \ell(\beta, \tilde{\beta}) = (\tilde{\beta} - \beta)^2, \ell(\lambda, \tilde{\lambda}) = (\tilde{\lambda} - \lambda)^2, \quad (20)$$

where $\tilde{\alpha}, \tilde{\theta}, \tilde{\beta}$ and $\tilde{\lambda}$ denote the estimated posterior means of α, θ, β and λ , respectively.

The generalized entropy (GE), an asymmetric loss function, is a simple generalization of the entropy loss with the shape parameter q being 1 and is given by

$$\ell(\omega, \tilde{\omega}) \propto \left(\frac{\tilde{\omega}}{\omega}\right)^q - q \ln\left(\frac{\tilde{\omega}}{\omega}\right) - 1, \quad q \neq 1, \quad (21)$$

where $\tilde{\omega}$ is an approximated estimation of ω given by

$$\tilde{\omega}_{GE} = [E_{\omega}(\omega^{-q})]^{-\frac{1}{q}}, \quad (22)$$

assuming that ω^{-q} exists and is finite and E_{ω} represents the anticipated value [34]. It should be emphasized that other loss functions may easily be substituted in the same way.

6. Optimization Criterion

There has been a lot of interest in identifying the best censoring scheme in recent years; see [35–39]. For values of n and m determined by the samples under a test, possible censoring schemes are all combinations of R_1, \dots, R_m . We are interested in selecting the best sample technique, as it entails identifying the progressive censoring scheme that provides the most information about the unknown parameters among all conceivable progressive censoring schemes. The first challenge is to determine a way to generate the unknown parameter information based on specific progressive censoring data, and the second is to compare two distinct information measures based on two different progressive censoring techniques. We, here, provide some of the optimality criteria as follows. We choose the censoring method that provides the most information about the unknown parameters. Table 1 provides a variety of commonly used measures in selecting the proper progressive censoring strategy, C_i .

Table 1. Some practical censoring plan optimum criteria.

Criterion	Method
C_1	Maximize trace $[I_{4 \times 4}(\cdot)]$
C_2	Minimize trace $[I_{4 \times 4}(\cdot)]^{-1}$
C_3	Minimize det $[I_{4 \times 4}(\cdot)]^{-1}$

We are interested in maximizing the observed Fisher information $I_{4 \times 4}(\cdot)$ for C_1 . Furthermore, for criteria C_2 and C_3 , we reduce the determinant and trace of $[I_{4 \times 4}(\cdot)]^{-1}$. Comparing multiple criteria is simple when dealing with a distribution with a single parameter; however, comparing the two Fisher information matrices becomes difficult for multiparameter distributions, because C_2 and C_3 are not scale invariant. As a result, the logarithm of the APIW distribution for \hat{t}_p is provided by

$$\log(\hat{t}_p) = \log \left\{ \frac{-1}{\beta} \log \left[1 - \frac{\log(1 + p(\alpha - 1))}{\log \alpha} \right] \right\}^{\frac{1}{\theta}}, \quad 0 < p < 1, \quad (23)$$

We apply the delta approach to (23) to obtain the approximated variance for $\log(\hat{t}_p)$ of the APIW distribution as

$$\text{Var}(\log(\hat{t}_p)) = [\nabla \log(\hat{t}_p)]^T I_{4 \times 4}^{-1}(\hat{\alpha}, \hat{\theta}, \hat{\beta}, \hat{\lambda}) [\nabla \log(\hat{t}_p)],$$

where

$$[\nabla \log(\hat{t}_p)]^T = \left[\frac{\partial}{\partial \alpha} \log(\hat{t}_p), \frac{\partial}{\partial \theta} \log(\hat{t}_p), \frac{\partial}{\partial \beta} \log(\hat{t}_p), \frac{\partial}{\partial \lambda} \log(\hat{t}_p) \right]_{(\alpha=\hat{\alpha}, \theta=\hat{\theta}, \beta=\hat{\beta}, \lambda=\hat{\lambda})}$$

The optimal progressive censoring corresponds to a maximum value of C_1 and a minimum value of $C_i, i = 1, 2, 3$.

7. Simulation

In this section, simulation experiments are carried out to evaluate the MLEs and Bayesian estimators' performances under the SLF and ELF in terms of their bias, mean square error (MSE), length of asymptotic CIs (LACI), and length of credible CIs (LCCI). The 95% CIs are generated using the asymptotic distribution of the MLEs. Two MLE bootstrap confidence intervals are additionally attained. The HPD is used to calculate the 95% credible intervals. Two schemes of progressive censoring are taken into consideration:

Scheme I: $R_1 = \dots = R_{m-1} = 0$, and $R_m = n - m$.

Scheme II: $R_2 = \dots = R_m = 0$ and $R_1 = n - m$.

For more information, see [40]. To choose the best strategy for the determinant and trace of the variance–covariance matrices, maximization of the principal diagonal elements of the Fisher information matrices, minimization of the determinant and trace of the variance–covariance matrix, and minimization of the variance in the logarithmic MLE of the p -th quantile, we used various optimization criteria. The following algorithm is used to carry out the estimation procedure:

1. Give the numbers n, m , and τ . The total sample size in complete case is $n = 100$ and 200 ; the censored sample size is $m = 75$ and 90 m when $n = 100$ and $m = 150$ and 185 when $n = 200$.
2. Give the parameters, $\alpha = 2, \beta = 2, \lambda = 1.6, \theta = 0.7$, and $\alpha = 0.6, \beta = 0.7, \lambda = 0.8, \theta = 1.4$.
3. Make a sample of size n of the randomness from the random variable t in Equation (1), then sort it. It is easy to create a random variable with the APIW distribution. If the uniform random variable U is drawn from the interval $[0, 1]$, then

$$z = \begin{cases} \left\{ \frac{-1}{\beta} \ln \left[\frac{\ln(1+(\alpha-1)U)}{\ln(\alpha)} \right] \right\}^{\frac{-1}{\theta}} & t < \tau \\ \tau - \tau^* + \left\{ \frac{-1}{\lambda} \ln \left[\frac{\ln(1+(\alpha-1)u)}{\ln(\alpha)} \right] \right\}^{\frac{-1}{\theta}} & t > \tau \end{cases}$$

4. To generate the adaptive progressive hybrid censored data for given n, m , and δ , we use the model in (7). The data can be thought of as:

$$z_{1:m:n} < z_{2:m:n} < \dots < z_{u:m:n} < \tau < z_{u+1:m:n} < \dots < z_{c:m:n} < \delta < z_{c+1:m:n} < z_{m:m:n}$$

5. To obtain the MLEs of the parameters, the nonlinear system is solved by using the Newton–Raphson method.
6. To obtain the Bayes estimation of the parameters, we obtain posterior samples from the MCMC algorithm.
7. Repeat Steps 3 through 6 for 1000 iterations.
8. Calculate the MLEs and Bayes parameter-related average values of bias, MSE, and LCI.
9. Calculate various parameter estimations and their confidence intervals.
10. Calculate the various optimization criteria.

Numerical simulation studies are provided in Tables 2–7 and Figures 2 and 3. Several conclusions can be drawn as follows.

- By increasing the censored sample sizes m , the bias, MSE, and LCI of the estimates for the two alternative censored methods decrease for fixed values of n and δ .

- By increasing the censored sample sizes δ , the bias, MSE, and LCI of the estimates for the two alternative censored methods decrease for fixed values of n and m .
- By increasing the censored sample sizes n , the bias, MSE, and LCI of the estimates for the two alternative censored methods decrease for fixed values of the sample sizes δ and m .
- Bayes estimations of the parameters under the two loss functions outperform the MLE in terms of bias and MSE for the scenarios under consideration.
- The bias and MSE of Bayes estimations of the parameters increase under the considered scenarios when we used negative weight for ELF.
- The HPDs of the unknown parameters outperform the CIs based on the MLEs with respect to ACIs and LCIs. In addition, we observe that the lengths of the bootstrap CIs are the shortest.

Table 2. Bias, MSE, LACI, LBCI, and LCCI with scheme 1 in case 1.

$\alpha = 2, \beta = 2, \lambda = 1.6, \theta = 0.7$			MLE					SELF			ELF $c = -1.25$			ELF $c = 1.25$			
n	τ, δ	m	Bias	MSE	LACI	LBPCI	LBTCI	Bias	MSE	LCCI	Bias	MSE	LCCI	Bias	MSE	LCCI	
100	2, 10	75	α	0.3263	0.3997	2.1248	0.0965	0.0947	-0.2602	0.3062	1.8368	-0.2458	0.2894	1.7968	-0.4100	0.5367	2.2491
			β	-1.3825	1.9408	0.6726	0.0302	0.0301	-0.5628	0.3439	0.6222	-0.5578	0.3379	0.6157	-0.6050	0.3980	0.6726
			λ	-0.3294	0.1665	0.9447	0.0406	0.0406	0.2863	0.1435	0.7539	0.6361	0.4416	0.7597	0.5878	0.3778	0.6959
		θ	0.8792	0.8010	0.6577	0.0301	0.0300	0.0546	0.0395	0.3513	0.0563	0.0097	0.3143	0.0389	0.0079	0.3100	
		90	α	0.1289	0.3267	2.1851	0.1033	0.1031	-0.0321	0.0331	0.7063	-0.0309	0.0329	0.7056	-0.0423	0.0351	0.7149
			β	-1.3616	1.8885	0.7287	0.0325	0.0327	-0.2817	0.1017	0.5869	-0.2789	0.0997	0.5807	-0.3071	0.1212	0.6382
	λ		-0.2883	0.1568	1.0653	0.0484	0.0484	0.1990	0.0575	0.5284	0.2005	0.0583	0.5303	0.1850	0.0506	0.4982	
	3.5, 18	75	θ	0.8403	0.7273	0.5710	0.0257	0.0253	0.0464	0.0241	0.3401	0.1281	0.0246	0.3411	0.1123	0.0198	0.3214
			α	1.2072	2.6719	4.3243	0.1911	0.1921	-0.1065	0.2610	1.8586	-0.0961	0.2537	1.8425	-0.2112	0.3636	2.1097
			β	-1.1084	1.2923	0.9910	0.0427	0.0432	-0.6102	0.4100	0.7449	-0.6050	0.4028	0.7387	-0.6526	0.4709	0.7836
		90	λ	0.4902	0.4963	1.9853	0.0901	0.0900	0.2893	0.2940	0.6019	0.9399	0.9551	1.0307	0.8542	0.7929	0.9038
			θ	0.6934	0.5379	0.9372	0.0414	0.0410	0.1369	0.0546	0.4633	0.1387	0.0560	0.4645	0.1201	0.0390	0.4486
α			1.3165	2.5775	3.6059	0.1659	0.1640	-0.0176	0.0289	0.6324	-0.0166	0.0288	0.6321	-0.0262	0.0298	0.6365	
200	150	β	-0.7230	0.9408	0.6724	0.0317	0.0317	-0.2981	0.1098	0.5521	-0.2953	0.1077	0.5458	-0.3236	0.1297	0.6095	
		λ	0.3408	0.2248	1.2936	0.0615	0.0615	0.2828	0.0994	0.5239	0.2850	0.1009	0.5256	0.2630	0.0863	0.4918	
		θ	0.5154	0.5080	0.4873	0.0216	0.0217	0.1171	0.0185	0.2572	0.1185	0.0188	0.2583	0.1050	0.0154	0.2438	
	185	α	0.4348	0.3536	1.5917	0.0687	0.0686	-0.3758	0.3113	1.5792	-0.3615	0.2930	1.5336	-0.5182	0.5351	1.9342	
		β	-1.4213	2.0289	0.3648	0.0164	0.0164	-0.5234	0.2866	0.4452	-0.5207	0.2836	0.4421	-0.5462	0.3127	0.4733	
		λ	-0.3996	0.1784	0.5367	0.0252	0.0251	0.3067	0.1470	0.4559	0.6752	0.4750	0.5652	0.6380	0.4234	0.5026	
300	2, 10	θ	0.8859	0.7991	0.4679	0.0216	0.0216	0.1493	0.0536	0.2133	0.0502	0.0055	0.2140	0.0411	0.0046	0.2089	
		α	0.3155	0.2366	1.4526	0.0660	0.0657	-0.0758	0.0305	0.6101	-0.0749	0.0303	0.6091	-0.0847	0.0331	0.6135	
		β	-1.4129	2.0049	0.3619	0.0153	0.0150	-0.3626	0.1448	0.4347	-0.3595	0.1423	0.4300	-0.3899	0.1677	0.4811	
	150	λ	-0.3822	0.1643	0.5300	0.0236	0.0241	0.2720	0.0855	0.4317	0.2737	0.0866	0.4337	0.2565	0.0760	0.4061	
		θ	0.8609	0.7536	0.4379	0.0193	0.0194	0.1081	0.0367	0.2046	0.1822	0.0372	0.2442	0.1678	0.0318	0.2363	
		α	0.4026	0.2824	1.4873	0.0552	0.0567	-0.1721	0.2094	1.5946	-0.1630	0.2006	1.5636	-0.2593	0.3096	1.8659	
450	150	β	-0.9198	1.4616	0.3652	0.0129	0.0128	-0.5550	0.3255	0.5263	-0.5521	0.3219	0.5242	-0.5792	0.3553	0.5561	
		λ	0.2994	0.1698	0.5294	0.0258	0.0255	0.1025	0.1085	0.4657	1.0326	1.1012	0.6591	0.9497	0.9289	0.5761	
		θ	0.7250	0.5558	0.6811	0.0314	0.0309	0.1129	0.0419	0.2710	0.1138	0.0197	0.2710	0.1047	0.0168	0.2699	
	185	α	0.4004	0.2704	1.0908	0.0425	0.0416	-0.0208	0.0234	0.5713	-0.0200	0.0233	0.5702	-0.0283	0.0244	0.5746	
		β	-0.2134	0.8482	0.3493	0.0109	0.0119	-0.3858	0.1599	0.3899	-0.3823	0.1570	0.3850	-0.4166	0.1867	0.4370	
		λ	0.2444	0.0915	0.5070	0.0219	0.0214	0.2369	0.0877	0.4251	0.3718	0.1509	0.4277	0.3447	0.1295	0.3881	
θ	0.6121	0.4589	0.3919	0.0174	0.0174	0.1670	0.0305	0.1924	0.1681	0.0309	0.1941	0.1565	0.0268	0.1823			

Table 3. Bias, MSE, LACI, LBCI, and LCCI with scheme 2 in case 1.

$\alpha = 2, \beta = 2, \lambda = 1.6, \theta = 0.7$			MLE				SELF			ELF $c = -1.25$			ELF $c = 1.25$				
n	τ, δ	m	Bias	MSE	LACI	LBPCI	LBTCI	Bias	MSE	LCCI	Bias	MSE	LCCI	Bias	MSE	LCCI	
100	2, 10	75	α	0.4624	0.5714	2.3474	0.1201	0.1205	-0.2218	0.2633	1.8289	-0.2105	0.2495	1.7748	-0.3394	0.4512	2.2924
			β	-1.3916	1.9624	0.6328	0.0319	0.0321	-0.5221	0.2971	0.6066	-0.5175	0.2918	0.5994	-0.5610	0.3430	0.6561
			λ	-0.3496	0.1814	0.9551	0.0496	0.0499	0.3042	0.1802	0.7407	0.4214	0.2150	0.7430	0.3813	0.1781	0.7106
			θ	1.0136	1.0650	0.7624	0.0425	0.0421	0.5739	0.3628	0.7099	0.5790	0.3695	0.7194	0.5214	0.2973	0.6187
	3.5, 18	90	α	0.5295	0.5660	2.4170	0.0765	0.0770	-0.2670	0.2579	1.7531	-0.2544	0.2632	1.6937	-0.4003	0.5122	2.3486
			β	-1.4030	1.9596	0.6540	0.0210	0.0206	-0.5261	0.2965	0.5475	-0.5217	0.2915	0.5396	-0.5636	0.3405	0.5825
			λ	-0.3740	0.1798	1.3282	0.0431	0.0425	0.4374	0.1800	0.7370	0.4412	0.2313	0.7418	0.4023	0.1940	0.6995
			θ	0.9861	1.0122	0.7831	0.0251	0.0249	0.5751	0.3640	0.6653	0.5798	0.3701	0.6715	0.5259	0.3017	0.6024
	2, 10	150	α	0.4601	0.5524	2.2904	0.1124	0.1125	-0.1171	0.2100	1.6852	-0.1090	0.2033	1.6784	-0.1962	0.2943	1.8787
			β	-1.1979	1.4732	0.7660	0.0243	0.0246	-0.5480	0.2832	0.5410	-0.5432	0.3140	0.5365	-0.5884	0.3679	0.5782
			λ	0.2992	0.1722	0.9408	0.0444	0.0438	0.6883	0.1526	0.9032	0.6950	0.5369	0.9078	0.6229	0.4307	0.8363
			θ	0.8304	0.7148	0.6230	0.0188	0.0188	0.4609	0.2310	0.5173	0.4644	0.2346	0.5208	0.4256	0.1959	0.4685
3.5, 18	90	α	0.5074	0.5339	2.2637	0.0712	0.0716	-0.0227	0.0252	0.5770	-0.0219	0.0251	0.5750	-0.0300	0.0263	0.5855	
		β	-1.1968	1.4705	0.7666	0.0240	0.0241	-0.2323	0.0708	0.5093	-0.2303	0.0697	0.5044	-0.2500	0.0822	0.5517	
		λ	0.2893	0.1719	0.9128	0.0292	0.0298	0.2058	0.0585	0.4962	0.2073	0.0593	0.4986	0.1917	0.0513	0.4786	
		θ	0.8319	0.7175	0.6251	0.0204	0.0205	0.2410	0.0649	0.3098	0.2432	0.0661	0.3141	0.2204	0.0542	0.2832	
200	2, 10	150	α	0.4968	0.4611	1.8157	0.0594	0.0594	-0.4156	0.3868	1.7927	-0.3966	0.3570	1.7173	-0.6259	0.8221	2.3011
			β	-1.4238	2.0385	0.4153	0.0131	0.0131	-0.5478	0.3116	0.4096	-0.5445	0.3078	0.4078	-0.5748	0.3434	0.4355
			λ	-0.4137	0.1950	0.6052	0.0186	0.0184	0.4819	0.1826	0.5607	0.4852	0.2608	0.6138	0.4507	0.2244	0.5572
			θ	0.9426	0.9101	0.5755	0.0177	0.0173	0.5626	0.3389	0.5679	0.5658	0.3428	0.5713	0.5291	0.2985	0.5178
	3.5, 18	185	α	0.5008	0.4615	1.8002	0.0559	0.0563	-0.0585	0.0268	0.5996	-0.0576	0.0265	0.5964	-0.0668	0.0293	0.6242
			β	-1.4305	2.0554	0.3718	0.0120	0.0118	-0.3356	0.1240	0.4175	-0.3328	0.1220	0.4126	-0.3604	0.1433	0.4485
			λ	-0.4246	0.1920	0.5577	0.0183	0.0180	0.2406	0.0689	0.4180	0.2421	0.0698	0.4216	0.2260	0.0609	0.3888
			θ	0.9422	0.9061	0.5310	0.0167	0.0167	0.3808	0.1519	0.3178	0.3844	0.1548	0.3210	0.3454	0.1245	0.2811
	2, 10	150	α	0.4717	0.3649	1.7284	0.0574	0.0570	-0.2171	0.2180	1.5888	-0.2071	0.2071	1.5436	-0.3190	0.3637	2.0831
			β	-1.2425	1.5586	0.4772	0.0152	0.0148	-0.5556	0.3018	0.3705	-0.5524	0.3143	0.3680	-0.5818	0.3487	0.3847
			λ	0.1907	0.0833	0.8496	0.0252	0.0254	0.8361	0.0733	0.7085	0.8428	0.7451	0.7166	0.7687	0.6179	0.6418
			θ	0.7916	0.6403	0.4600	0.0147	0.0148	0.4665	0.2287	0.4089	0.4688	0.2310	0.4112	0.4432	0.2060	0.3780
3.5, 18	185	α	1.5699	0.3267	3.5125	0.1158	0.1166	-0.0406	0.0260	0.6052	-0.0398	0.0259	0.6024	-0.0483	0.0275	0.6174	
		β	-1.2296	1.5242	0.4327	0.0146	0.0144	-0.3682	0.1148	0.4297	-0.3648	0.1450	0.4256	-0.3978	0.1727	0.4703	
		λ	0.2119	0.0832	0.7672	0.0246	0.0253	0.3420	0.0513	0.4309	0.3447	0.1319	0.4339	0.3172	0.1117	0.4069	
		θ	0.7923	0.6399	0.4322	0.0138	0.0138	0.3183	0.1058	0.2583	0.3207	0.1074	0.2611	0.2952	0.0908	0.2329	

Table 4. Bias, MSE, LACI, LBCI, and LCCI with scheme 1 in case 2.

$\alpha = 0.6, \beta = 0.7, \lambda = 0.8, \theta = 1.4$			MLE				SELF			ELF $c = -1.25$			ELF $c = 1.25$				
n	τ, δ	m	Bias	MSE	LACI	LBPCI	LBTCI	Bias	MSE	LCCI	Bias	MSE	LCCI	Bias	MSE	LCCI	
100	0.6, 1.3	75	α	-0.5240	0.2747	0.1405	0.0022	0.0022	-0.1110	0.0206	0.3572	-0.1057	0.0196	0.3578	-0.1615	0.0343	0.3626
			β	-0.2999	0.0922	0.2150	0.0086	0.0083	-0.2663	0.0740	0.2063	-0.2641	0.0728	0.2066	-0.2856	0.0848	0.2194
			λ	0.0052	0.0093	0.4453	0.0177	0.0177	0.0200	0.0083	0.3513	0.0222	0.0084	0.3514	0.0008	0.0081	0.3521
			θ	0.2704	0.0885	0.4687	0.0205	0.0209	0.0732	0.0181	0.4339	0.0750	0.0184	0.4339	0.0574	0.0156	0.4351
	3.5, 1.3	90	α	-0.4477	0.2017	0.0482	0.0061	0.0061	-0.0486	0.0051	0.2093	-0.0475	0.0050	0.2086	-0.0591	0.0064	0.2144
			β	-0.2570	0.0691	0.1885	0.0098	0.0100	-0.2212	0.0516	0.2015	-0.2192	0.0507	0.1992	-0.2383	0.0599	0.2146
			λ	0.0945	0.0082	0.3785	0.0203	0.0203	0.0211	0.0056	0.2751	0.0220	0.0056	0.2750	0.0129	0.0055	0.2783
			θ	0.1132	0.0271	0.4865	0.0212	0.0202	0.0282	0.0104	0.3800	0.0289	0.0105	0.3799	0.0218	0.0099	0.3766

Table 4. Cont.

$\alpha = 0.6, \beta = 0.7, \lambda = 0.8, \theta = 1.4$			MLE				SELF			ELF c = -1.25			ELF c = 1.25				
n	τ, δ	m	Bias	MSE	LACI	LBPCI	LBTCI	Bias	MSE	LCCI	Bias	MSE	LCCI	Bias	MSE	LCCI	
200	0.8, 1.5	75	α	-0.5342	0.2856	0.1804	0.0019	0.0019	-0.0505	0.0780	0.5301	-0.0449	0.0815	0.5360	-0.0990	0.0461	0.5708
		β	-0.1805	0.0375	0.3960	0.0082	0.0082	-0.2799	0.0528	0.2684	-0.2775	0.0820	0.2668	-0.3001	0.0953	0.2745	
		λ	0.3202	0.1212	0.8482	0.0163	0.0161	0.1205	0.0312	0.4830	0.1232	0.0318	0.4820	0.0957	0.0263	0.4838	
		θ	0.2652	0.0891	0.5336	0.0169	0.0173	0.2184	0.0708	0.5753	0.2206	0.0720	0.5783	0.1984	0.0606	0.5573	
	90	α	-0.5122	0.2644	0.0611	0.0056	0.0057	-0.0334	0.0081	0.2086	-0.0323	0.0037	0.2086	-0.0431	0.0047	0.2083	
		β	-0.0762	0.0160	0.2736	0.0124	0.0127	-0.2306	0.0506	0.1998	-0.2285	0.0548	0.1982	-0.2477	0.0644	0.2139	
		λ	0.2548	0.1135	0.5362	0.0258	0.0258	0.1086	0.0193	0.3247	0.1096	0.0195	0.3253	0.0994	0.0171	0.3165	
		θ	0.0780	0.0246	0.5238	0.0167	0.0165	0.0940	0.0189	0.3921	0.0949	0.0191	0.3931	0.0858	0.0169	0.3768	
	150	α	-0.5503	0.3029	0.1706	0.0008	0.0008	-0.1207	0.0200	0.2611	-0.1166	0.0189	0.2599	-0.1571	0.0303	0.2621	
		β	-0.2948	0.0882	0.1971	0.0044	0.0044	-0.2850	0.0833	0.1726	-0.2835	0.0824	0.1715	-0.2982	0.0911	0.1726	
		λ	0.0074	0.0524	0.4162	0.0089	0.0090	0.0116	0.0057	0.2784	0.0131	0.0057	0.2773	-0.0023	0.0057	0.2796	
		θ	0.3037	0.1005	0.3554	0.0109	0.0109	0.0952	0.0189	0.3662	0.0964	0.0192	0.3668	0.0848	0.0166	0.3599	
185	α	-0.4551	0.2090	0.0259	0.0055	0.0055	-0.0324	0.0051	0.0733	-0.0320	0.0014	0.0732	-0.0362	0.0017	0.0757		
	β	-0.1976	0.0416	0.1409	0.0063	0.0062	-0.2553	0.0663	0.1278	-0.2536	0.0654	0.1269	-0.2687	0.0734	0.1345		
	λ	0.2050	0.0053	0.2817	0.0136	0.0137	-0.0202	0.0031	0.1965	-0.0193	0.0031	0.1962	-0.0277	0.0036	0.2046		
	θ	0.0212	0.0074	0.3277	0.0110	0.0110	0.0694	0.0068	0.2919	0.0701	0.0108	0.2922	0.0638	0.0097	0.2890		
150	α	-0.5588	0.3131	0.1154	0.0054	0.0054	-0.0395	0.2423	0.8773	-0.0266	0.4469	0.8702	-0.1289	0.1863	0.8322		
	β	0.0655	0.0179	0.4586	0.0212	0.0213	-0.0833	0.0109	0.3543	-0.2813	0.0898	0.4482	-0.3004	0.1008	0.4510		
	λ	0.8364	0.7483	0.8673	0.0399	0.0395	0.1402	0.0585	0.8695	0.1427	0.0593	0.8679	0.1171	0.0513	0.8502		
	θ	-0.0539	0.0232	0.5586	0.0267	0.0267	0.2571	0.0194	0.5065	0.2587	0.0953	0.6570	0.2422	0.0843	0.6294		
185	α	-0.5559	0.3091	0.0279	0.0010	0.0010	-0.0218	0.0594	0.0401	-0.0214	0.0006	0.0400	-0.0256	0.0008	0.0424		
	β	-0.0517	0.0133	0.1916	0.0058	0.0059	-0.3182	0.0102	0.1041	-0.3169	0.1011	0.1039	-0.3293	0.1092	0.1060		
	λ	0.3258	0.1151	0.3711	0.0120	0.0120	0.0068	0.0027	0.1917	0.0082	0.0027	0.1917	-0.0056	0.0027	0.1963		
	θ	0.0429	0.0193	0.3909	0.0122	0.0122	0.0290	0.0190	0.3326	0.2915	0.0924	0.3330	0.2772	0.0838	0.3223		

Table 5. Bias, MSE, LACI, LBCI, and LCCI with scheme 2 in case 2.

$\alpha = 0.6, \beta = 0.7, \lambda = 0.8, \theta = 1.4$			MLE				SELF			ELF c = -1.25			ELF c = 1.25				
n	τ, δ	m	Bias	MSE	LACI	LBPCI	LBTCI	Bias	MSE	LCCI	Bias	MSE	LCCI	Bias	MSE	LCCI	
100	0.6, 1.3	75	α	-0.6947	0.4827	0.6661	0.0015	0.0015	-0.0664	0.0453	0.6946	-0.0642	0.0425	0.6770	-0.0647	0.0439	0.6710
		β	-0.2988	0.0909	0.1620	0.0357	0.0364	-0.3179	0.0906	0.3675	-0.3146	0.0902	0.3623	-0.2934	0.0901	0.3400	
		λ	-0.0135	0.0093	0.3837	0.0865	0.0825	-0.1626	0.0090	0.4409	-0.1595	0.0383	0.4353	-0.1589	0.0085	0.4282	
		θ	0.6069	0.3911	0.6072	0.1366	0.1323	0.3054	0.1329	0.8110	0.3081	0.1351	0.8130	0.2801	0.1132	0.7882	
	90	α	-0.6950	0.4830	0.6182	0.0030	0.0026	-0.0019	0.0411	0.0079	-0.0018	0.0009	0.0073	-0.0013	0.0250	0.0013	
		β	-0.2316	0.0718	0.1572	0.0346	0.0345	-0.2133	0.0712	0.2092	-0.2033	0.0691	0.2085	-0.2104	0.0714	0.2008	
		λ	-0.0455	0.0074	0.3045	0.1013	0.1010	-0.1628	0.0068	0.2907	-0.1591	0.0063	0.2908	-0.1595	0.0062	0.2801	
		θ	0.5775	0.3441	0.4313	0.1406	0.1400	0.2543	0.1010	0.6715	0.2581	0.1036	0.6755	0.2197	0.0787	0.6262	
	75	α	-0.5943	0.3532	0.4910	0.0016	0.0016	0.0584	0.0421	0.5866	0.0626	0.2306	0.5715	0.0198	0.0370	0.5360	
		β	-0.1657	0.0340	0.3209	0.0554	0.0552	-0.3376	0.0291	0.3288	-0.3337	0.1182	0.3236	-0.3267	0.0214	0.3036	
		λ	0.3389	0.0090	0.6334	0.1018	0.1016	-0.0575	0.0079	0.6268	-0.0537	0.0270	0.6184	-0.0491	0.0063	0.6080	
		θ	0.4365	0.2228	0.7154	0.1166	0.1156	0.3705	0.1272	0.6530	0.3735	0.1275	0.6568	0.3411	0.1146	0.6132	
90	α	-0.5941	0.3529	0.4108	0.0021	0.0020	0.0677	0.0412	0.4675	0.2800	1.1994	3.1964	0.0612	0.0326	0.4268		
	β	-0.1699	0.0315	0.3000	0.0550	0.0556	-0.3659	0.0139	0.2499	-0.3616	0.1357	0.2466	-0.3399	0.0167	0.2317		
	λ	0.3172	0.0012	0.5363	0.1035	0.1007	-0.1149	0.0012	0.4647	-0.1103	0.0273	0.4588	-0.1056	0.0015	0.4510		
	θ	0.4460	0.2231	0.6214	0.1171	0.1176	0.4044	0.1010	0.6072	0.4082	0.2037	0.4728	0.3688	0.0917	0.5640		

Table 5. Cont.

$\alpha = 0.6, \beta = 0.7,$ $\lambda = 0.8, \theta = 1.4$			MLE					SELF			ELF $c = -1.25$			ELF $c = 1.25$			
n	τ, δ	m	Bias	MSE	LACI	LBPCI	LBTCI	Bias	MSE	LCCI	Bias	MSE	LCCI	Bias	MSE	LCCI	
200	0.6, 1.3	150	α	-0.5964	0.3557	0.0039	0.0007	0.0006	-0.0005	0.0011	0.0031	-0.0005	0.0001	0.0031	-0.0006	0.0001	0.0031
			β	-0.2865	0.0831	0.1284	0.0229	0.0226	-0.2953	0.0793	0.1128	-0.2927	0.0914	0.2801	-0.2315	0.0658	0.1030
			λ	0.0095	0.0029	0.2102	0.0385	0.0364	-0.0012	0.0023	0.2034	-0.1163	0.0221	0.3416	-0.0014	0.0229	0.2036
		θ	0.5218	0.2802	0.3539	0.0606	0.0602	0.2415	0.0756	0.3044	0.2434	0.0767	0.4460	0.2244	0.0656	0.3042	
		185	α	-0.4695	0.3248	0.0037	0.0006	0.0005	-0.0043	0.0005	0.0025	-0.0034	0.0001	0.0242	-0.0041	0.0001	0.0025
			β	-0.2321	0.0810	0.1210	0.0242	0.0237	-0.2402	0.0692	0.1028	-0.2840	0.0817	0.2766	-0.2143	0.0519	0.1030
	λ		-0.0157	0.0027	0.2039	0.0426	0.0427	-0.2624	0.0022	0.1931	-0.2583	0.0804	0.4188	-0.2594	0.0022	0.1845	
	0.8, 1.5	150	θ	0.4604	0.2372	0.3384	0.0614	0.0612	0.4370	0.0622	0.2613	0.4406	0.2277	0.6197	0.4028	0.0619	0.2546
			α	-0.4693	0.3248	0.0038	0.0015	0.0015	0.1955	0.0013	0.0025	0.2065	1.2956	0.6617	0.0749	0.0123	0.0022
			β	-0.1932	0.0418	0.2645	0.0396	0.0393	-0.3756	0.0415	0.2195	-0.3719	0.1504	0.2897	-0.3403	0.0402	0.2132
		185	λ	0.0029	0.0020	0.2051	0.0806	0.0806	-0.1400	0.0020	0.1499	-0.1361	0.0399	0.4920	-0.1373	0.0016	0.1542
			θ	0.4195	0.2318	0.2620	0.0925	0.0930	0.5393	0.0720	0.2279	0.5434	0.3563	0.7944	0.4997	0.0630	0.2175
α			-0.3692	0.3048	0.0010	0.0011	0.0011	-0.0521	0.0004	0.0023	-0.0475	0.0905	0.6529	-0.0471	0.0047	0.1700	
β	-0.1882	0.0415	0.2329	0.0269	0.0268	-0.3041	0.0315	0.1925	-0.2409	0.1746	0.2626	-0.2944	0.0302	0.1832			
λ	0.0025	0.0018	0.1943	0.0503	0.0487	-0.1873	0.0014	0.1476	-0.1829	0.0494	0.4703	-0.1822	0.0014	0.1490			
θ	0.3560	0.2033	0.2150	0.0605	0.0613	0.4586	0.0519	0.1597	0.5908	0.4063	0.9854	0.4154	0.0434	0.1486			

Table 6. Optimization criterion with different schemes and cases.

Case	n	Scheme τ, δ	1			2				
			m	C1	C2	C3	C1	C2	C3	
1	100	2, 10	75	8.9523	3.860×10^{-5}	758.8371	7.9682	4.248×10^{-5}	679.5787	
			90	7.1809	6.359×10^{-6}	766.3755	7.4242	3.969×10^{-5}	769.7234	
		3.5, 18	75	4.7788	3.619×10^{-5}	765.5838	7.6918	1.605×10^{-5}	690.4161	
			90	3.4849	3.370×10^{-6}	778.3303	7.4603	1.503×10^{-5}	777.0073	
		200	2, 10	150	3.4486	7.767×10^{-7}	1616.8601	4.0384	1.900×10^{-6}	1306.3727
				185	3.1947	5.709×10^{-7}	1645.6376	3.3101	8.940×10^{-7}	1619.5660
	3.5, 18		150	3.1965	6.971×10^{-7}	1731.1179	4.0136	1.163×10^{-6}	1561.8482	
			185	2.9820	2.556×10^{-7}	1735.4056	3.0424	2.965×10^{-7}	1607.0823	
	2	100	2, 10	75	7.9682	4.248×10^{-4}	679.5787	0.3979	3.289×10^{-10}	6985.3086
				90	7.4242	6.873×10^{-4}	769.7234	0.3933	2.030×10^{-10}	72883.3960
			3.5, 18	75	6.9178	1.605×10^{-4}	690.4161	0.3869	9.505×10^{-10}	104,014.7057
				90	6.6033	1.503×10^{-4}	779.0073	0.3773	1.246×10^{-10}	117,561.7575
200			2, 10	150	4.0384	1.900×10^{-6}	1306.3727	0.3284	3.465×10^{-11}	410,484.9941
				185	3.3101	8.940×10^{-6}	1619.5660	0.3244	1.216×10^{-12}	214,081.1995
		3.5, 18	150	3.3625	1.163×10^{-6}	1761.8482	0.2731	1.068×10^{-11}	140,775.1960	
			185	3.0424	2.965×10^{-6}	1807.0823	0.2412	1.554×10^{-12}	178,754.3036	

Table 7. MLE, SE, and different measures of goodness of fit.

	Estimates	SE	AIC	CAIC	BIC	HQIC	CVM	AD	KS	PVKS
α	26.9222	74.2080	121.2031	121.6099	127.6326	123.7319	0.0997	0.5233	0.0976	0.5859
β	235.4901	222.2091								
λ	6.4564	0.6125								




Figure 2. Heatmap for MSE with scheme 1 in case 1.




Figure 3. Heatmap for MSE with scheme 1 in case 2.

8. A Real-Data Application

We use examination data from [41] to illustrate the practical application of the proposed model. The following information represents the strength measured in GPA for single carbon fibers with gauge lengths of 10 mm and sample size of 63: 1.901, 2.132, 2.203, 2.228, 2.257, 2.350, 2.361, 2.396, 2.397, 2.445, 2.454, 2.474, 2.518, 2.522, 2.525, 2.532, 2.575, 2.614, 2.616, 2.618, 2.624, 2.659, 2.675, 2.738, 2.740, 2.856, 2.917, 2.928, 2.937, 2.937, 2.977, 2.996, 3.030, 3.125, 3.139, 3.145, 3.220, 3.223, 3.235, 3.243, 3.264, 3.272, 3.294, 3.332, 3.346, 3.377, 3.408, 3.435, 3.493, 3.501, 3.537, 3.554, 3.562, 3.628, 3.852, 3.871, 3.886, 3.971, 4.024, 4.027, 4.225, 4.395, and 5.020. Here, we use the modified Kolmogorov–Smirnov as a method for the goodness-of-fit test as follows:

The computational formula for the modified Kolmogorov–Smirnov statistic is then given by

$$D_{m:n} = \max(D_{m:n}^+, D_{m:n}^-),$$

where

$$D_{m:n}^+ = \max_i (\omega_{i:m:n} - F(z_{i:m:n}; \hat{\alpha}, \hat{\beta}, \hat{\theta}, \hat{\lambda}))$$

$$D_{m:n}^- = \max_i (F(z_{i:m:n}; \hat{\alpha}, \hat{\beta}, \hat{\theta}, \hat{\lambda}) - \omega_{i-1:m:n}),$$

$$\omega_{i:m:n} = 1 - \prod_{j=m-i+1}^m \frac{j + R_{m-j+1} + \dots + R_m}{j + 1 + R_{m-j+1} + \dots + R_m}.$$

Ref. [42] proposed a general-purpose goodness-of-fit test by first estimating the unknown parameters of the hypothesized distribution, then transforming the data to normality, and then testing the goodness of fit of the transformed data to normality. Then, along the lines of [42], the proposed test procedure is as follows:

1. Find the maximum likelihood estimate of the unknown parameter $\alpha, \beta, \theta, \lambda$, denoted by $\hat{\alpha}, \hat{\beta}, \hat{\theta}, \hat{\lambda}$, under the hypothesized model and calculate $v_{i:m:n} = F(z_{i:m:n}; \hat{\alpha}, \hat{\beta}, \hat{\theta}, \hat{\lambda})$ for $i = 1, \dots, m$.
2. Generate $y_{i:m:n}$ as $F^{-1}(v_{i:m:n})$ for $i = 1, \dots, m$.
3. Considering $y_{1:m:n}, \dots, y_{m:m:n}$ as a progressively Type-II censored data from an APIW distribution with $\alpha, \beta, \theta, \lambda$ and calculate the maximum likelihood estimates $\hat{\alpha}, \hat{\beta}, \hat{\theta}, \hat{\lambda}$.
4. Calculate $u_{i:m:n} = \Phi(y_{i:m:n})$ for $i = 1, \dots, m$.
5. Calculate $D_{m:n}$
6. Reject the null hypothesis at significance level δ if the test statistic exceeds the upper tail significance points.

For more information about the p -value for KS test for SSPALT samples, see [42–46].

In Table 7, we provide the MLEs with their standard errors (SEs) for the APIW parameters and different measures of goodness of fit as Akaike information criterion (AIC), Bayesian information criterion (BIC), corrected Akaike information criterion (CAIC), Hannan–Quinn information criterion (HQIC), Kolmogorov–Smirnov (KS) test and its p -value (PVKS), Anderson–Darling (AD), and Cramèr–von Mises (CVM).

The empirical CDF and its CDF fitted (left panel), and the histogram of the data and its fitted density function to the single carbon fiber data (right panel) are displayed in the top of Figure 4. Further, a graph of the PP plot (left panel) and QQ plot (right panel) of the APIW distribution is shown in bottom Figure 4.




Figure 4. Fitting plot of APIW distribution of single carbon fibers.

Figure 5 shows the profile log-likelihood function plots for the parameters of the APIW distribution. Figure 6 displays contour plots of the log-likelihood function for the APIW parameters, indicating that the MLEs can be uniquely estimated.




Figure 5. Graphs of profile log-likelihood function for the parameters for the APIW model.




Figure 6. Contour plots of the log-likelihood function for the parameters for the APIW model.

Numerical results for the single carbon fiber study are provided in Table 8. Table 9 contains the estimations based on the censored data. For a given fixed scheme, we observe

that Bayes estimates of the unknown parameters are close to the MLEs. Table 10 discusses the estimation of τ , which is given by equating $F_1(\tau) = F_2(\tau^*)$. Table 11 discusses different optimality measures for the MLE based on different schemes, illustrating that the proposed technique is quite satisfactory.

Table 8. The single carbon fiber study data based on SSPALT when $\tau = 3$ and $\delta = 3.8$.

Scheme	Before τ								After τ					
I	1.901	2.132	2.203	2.228	2.257	2.350	2.361	2.396	3.030	3.125	3.139	3.145	3.220	3.223
	2.397	2.445	2.454	2.474	2.518	2.522	2.525	2.532						
	2.575	2.614	2.616	2.618	2.624	2.659	2.675	2.738						
	2.740	2.856	2.917	2.928	2.937	2.937	2.977	2.996						
II	1.901	2.132	2.203	2.257	2.350	2.361	2.396	2.397	3.030	3.125	3.139	3.145	3.220	3.235
	2.445	2.454	2.474	2.518	2.522	2.575	2.614	2.616						
	2.618	2.624	2.659	2.675	2.738	2.740	2.917	2.928						
				2.937	2.937	2.977								
								3.971	4.024	4.225	4.395	5.020		

Table 9. The MLE and its SE and Bayesian and its SD with confidence intervals.

<i>m</i>		MLE				Bayesian				
		Estimates	SE	Lower	Upper	Estimates	SD	Lower	Upper	
Complete	63	α	71.8806	22.7927	27.2070	116.5542	75.6191	18.5540	38.7075	111.2613
		β	162.0261	66.5356	31.6163	292.4359	190.3651	30.1964	135.8759	251.3891
		λ	229.3353	91.8064	49.3948	409.2759	269.1464	40.5281	189.3768	340.9331
		θ	7.4069	0.4585	6.5082	8.3056	7.5259	0.2519	7.0329	7.9940
I	50	α	156.2681	73.6006	12.0110	300.5253	167.0087	52.8410	71.6085	266.4320
		β	200.3566	99.6606	5.0219	395.6913	214.5642	42.8232	134.4151	288.3952
		λ	287.9273	113.3318	65.7969	510.0577	302.9877	57.5639	207.9282	407.4394
		θ	7.8263	0.6827	6.4882	9.1643	7.8582	0.3078	7.2443	8.4240
II	50	α	218.2157	108.2157	6.1130	430.3184	226.5631	77.5768	95.4818	369.4948
		β	270.8559	93.4996	87.5967	454.1151	279.5810	48.5672	176.8138	367.8409
		λ	377.6384	117.2506	147.8272	607.4497	389.1168	65.6272	248.3191	499.9688
		θ	8.1569	0.6178	6.9461	9.3678	8.1828	0.3113	7.6173	8.8183

Table 10. Estimated τ .

Scheme	MLE	Bayes
complete	3.14407	3.14130
I	3.14227	3.13465
II	3.12476	3.12368

Table 11. Optimality measures.

	Complete	I	II
C1	15,642.68	74,500.21	124,729.4
C2	26,415,630	1.24×10^9	9.09×10^9
C3	36.46753	35.4808	29.99117

9. Conclusions

It is known that in life-testing and reliability trials, many data may exhibit different shapes and are censored due to time and cost constraints. Thus, accelerated life tests are

commonly used to explore the lifetime of reliable items by subjecting them to elevated stress levels of stressors that could cause early failures. This observation motivated us to investigate the step stress partially accelerated life testing model with samples from the APIW distribution under the adaptive type II progressively hybrid censoring. We considered statistical inferences of the unknown model parameters of the APIW distribution from both likelihood and Bayesian perspectives. We first considered the maximum likelihood estimates for the unknown model parameters and used these estimates to construct two types of approximate confidence intervals of the distributional parameters. We then conducted Bayesian inference for the unknown parameters with non-informative and informative priors under the symmetric and asymmetric loss functions. Moreover, we analyzed three different probable optimum test techniques for the proposed model under different optimal criteria. Numerical results from both simulations and a real-data application illustrated that the performance of the proposed method is quite satisfactory for estimating the APIW parameters under different sampling schemes. We may, thus, conclude that the proposed model has great potential for analyzing censored data under the AT-II PHCS in the study of life testing and reliability analyses.

Author Contributions: R.A., E.M.A., Q.H., and H.R. have contributed equally. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2022R50), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data used to support the findings of this study are included within the article.

Acknowledgments: Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2022R50), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Fisher Information Matrix

The Fisher information matrix is a fundamental statistical construct that describes how much information data offer on a variable that is unknown. It can be used to calculate the variance in an estimator as well as the asymptotic behavior of maximum-likelihood estimations. The inverse of the Fisher information matrix is an estimator of the asymptotic covariance matrix. The Fisher information matrix is computed by taking the expected values of the negative second-partial and mixed-partial derivatives of the log-likelihood function with respect to α , θ , β and λ . It is further explained below.

$$I_{4 \times 4} = -E \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}, \tag{A1}$$

where $a_{11} = E\left(\frac{\partial^2 \ell}{\partial \alpha^2}\right)$, $a_{12} = a_{21} = E\left(\frac{\partial^2 \ell}{\partial \alpha \partial \theta}\right)$, $a_{13} = a_{31} = E\left(\frac{\partial^2 \ell}{\partial \alpha \partial \beta}\right)$, $a_{14} = a_{41} = E\left(\frac{\partial^2 \ell}{\partial \alpha \partial \lambda}\right)$, $a_{22} = E\left(\frac{\partial^2 \ell}{\partial \theta^2}\right)$, $a_{33} = E\left(\frac{\partial^2 \ell}{\partial \beta^2}\right)$, $a_{32} = a_{23} = E\left(\frac{\partial^2 \ell}{\partial \theta \partial \beta}\right)$, $a_{44} = E\left(\frac{\partial^2 \ell}{\partial \lambda^2}\right)$, $a_{42} = a_{24} = E\left(\frac{\partial^2 \ell}{\partial \theta \partial \lambda}\right)$, and $a_{34} = a_{34} = E\left(\frac{\partial^2 \ell}{\partial \beta \partial \lambda}\right)$.

The relevant matrices' elements are computed. As a result, the variance–covariance matrix for MLEs can be constructed as follows:

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \alpha^2} &= -\frac{(m_u+m)}{(\alpha \log \alpha)^2} + \frac{(m_u+m+\varepsilon_m)}{(\alpha-1)^2} - \frac{1}{\alpha^2} \sum_{i=1}^{m_u} e^{-\beta z_i^{-\theta}} + \frac{1}{\alpha^2} \sum_{i=1}^{m_u} \varepsilon_i + \frac{1}{(\alpha-1)^2} \sum_{i=1}^{m_u} \varepsilon_i \\
 &\quad - \left[\frac{1}{\alpha} \sum_{i=1}^{m_u} \varepsilon_i \alpha e^{-\beta z_i^{-\theta}} - 1 + (\log(\alpha))^2 \sum_{i=1}^{m_u} \varepsilon_i \alpha e^{-\beta z_i^{-\theta}} - 1 \right] - \frac{1}{\alpha^2} \sum_{i=m_u+1}^m \varepsilon_i e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} \\
 &\quad - \frac{1}{\alpha^2} \sum_{i=m_u+1}^m (\varepsilon_i) + \frac{1}{(\alpha-1)^2} \sum_{i=m_u+1}^m (\varepsilon_i) \\
 &\quad + \left[\frac{1}{\alpha} \sum_{i=m_u+1}^m \varepsilon_i \frac{\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1}{(1-\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1)} \right. \\
 &\quad + \log(\alpha) \sum_{i=m_u+1}^m \varepsilon_i (e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} \\
 &\quad \left. - 1) \left(\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 2 \right) \left(\frac{(1-\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1) + 1}{(1-\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1)^2} \right) \right] - \frac{\varepsilon_m}{\alpha^2} \\
 &\quad + \left[\frac{1}{\alpha} \sum_{i=m_u+1}^m \varepsilon_i \frac{\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1}{(1-\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1)} \right. \\
 &\quad + \log(\alpha) \sum_{i=m_u+1}^m \varepsilon_i (e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} \\
 &\quad \left. - 1) \left(\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 2 \right) \left(\frac{(1-\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1) + 1}{(1-\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1)^2} \right) \right], \tag{A2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \alpha \partial \theta} &= \frac{1}{\alpha} \sum_{i=1}^{m_u} e^{-\beta z_i^{-\theta}} \log(\beta z_i) - \log(\alpha) \sum_{i=1}^{m_u} \varepsilon_i e^{-\beta z_i^{-\theta}} \log(\beta z_i) \alpha e^{-\beta z_i^{-\theta}} - 1 + \frac{1}{\alpha} \sum_{i=m_u+1}^m \varepsilon_i \log(\beta[\tau + \\
 &\quad \lambda(z_i - \tau)]) e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} + \log(\alpha) \sum_{i=m_u+1}^m \varepsilon_i \left(\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1 \right) e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} \log(\beta[\tau + \\
 &\quad \lambda(z_i - \tau)]) \left(\frac{(1-\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1) + (\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1)}{(1-\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1)^2} \right) + \\
 &\quad \varepsilon_m \log(\alpha) \left(\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1 \right) e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} \log(\beta[\tau + \lambda(z_m - \\
 &\quad \tau)]) \left(\frac{(1-\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1) + (\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1)}{(1-\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1)^2} \right), \tag{A3}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \alpha \partial \beta} &= -\frac{1}{\alpha} \sum_{i=1}^{m_u} e^{-\beta z_i^{-\theta}} z_i^{-\theta} - \log(\alpha) \sum_{i=1}^{m_u} \varepsilon_i \alpha e^{-\beta z_i^{-\theta}} - 1 e^{-\beta z_i^{-\theta}} z_i^{-\theta} - \frac{1}{\alpha} \sum_{i=m_u+1}^m \varepsilon_i [\tau + \lambda(z_i - \tau)]^{-\theta} e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} \\
 &\quad + \log(\alpha) \sum_{i=m_u+1}^m \varepsilon_i \left(\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1 \right) [\tau \\
 &\quad + \lambda(z_i - \tau)]^{-\theta} e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} \left[\frac{(\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1) + 1}{(1-\alpha e^{-\beta[\tau+\lambda(z_i-\tau)]^{-\theta}} - 1)^2} \right] \\
 &\quad + \varepsilon_m \log(\alpha) \left(\alpha e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}} - 1 \right) [\tau + \lambda(z_m - \tau)]^{-\theta} e^{-\beta[\tau+\lambda(z_m-\tau)]^{-\theta}}, \tag{A4}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \alpha \partial \lambda} = & + \frac{1}{\alpha} \sum_{i=m_u+1}^m \varepsilon_i (z_i - \tau) e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \\
 & + \log(\alpha) \sum_{i=m_u+1}^m \varepsilon_i e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \beta(z_i - \tau) [\tau \\
 & + \lambda(z_i - \tau)]^{-\theta-1} \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \left(\frac{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1\right) + 1}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1\right)^2} \right) \\
 & + \varepsilon_m \log(\alpha) e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \beta(z_m - \tau) [\tau \\
 & + \lambda(z_m - \tau)]^{-\theta-1} \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \left(\frac{\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 + 1}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1\right)^2} \right), \tag{A5}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \theta^2} = & \beta \sum_{i=1}^{m_u} z_i^{-\theta} (\log(z_i))^2 \\
 & + \log(\alpha) \beta \sum_{i=1}^{m_u} \left(z_i^{-\theta} (\log(z_i))^2 e^{-\beta z_i^{-\theta}} - z_i^{-2\theta} \log(z_i) e^{-\beta z_i^{-\theta}} \right) \\
 & + \beta \sum_{i=1}^{m_u} \varepsilon_i e^{-\beta z_i^{-\theta}} z_i^{-\theta} \log(z_i) \left(e^{-\beta z_i^{-\theta}} - 1 \right) \alpha e^{-\beta z_i^{-\theta}} - 2 + \beta \sum_{i=m_u+1}^m [\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] \\
 & - \sum_{i=m_u+1}^m \log[\tau + \lambda(z_i - \tau)] + \log(\alpha) \sum_{i=m_u+1}^m e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \beta[\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] \\
 & + \sum_{i=m_u+1}^m \beta \varepsilon_i [\log[\tau + \lambda(z_i - \tau)]]^2 [\tau + \lambda(z_i - \tau)]^{-\theta} \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 \right) \\
 & + \left\{ \frac{\left[\left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \left[1 + \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) - [\tau + \lambda(z_i - \tau)]^{-\theta} + (\alpha e)^{-1} + \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right] \right]}{1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1} \right\} \\
 & + \beta \varepsilon_m (\log[\tau + \lambda(z_m - \tau)])^2 [\tau + \lambda(z_m - \tau)]^{-\theta} e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right) \\
 & + \left\{ \frac{\left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \left[1 + \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) - [\tau + \lambda(z_m - \tau)]^{-\theta} + (\alpha e)^{-1} + \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right]}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)} \right\}, \tag{A6}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \theta \partial \beta} = & \sum_{i=1}^{m_u} z_i^{-\theta} \log(z_i) + \log(\alpha) \sum_{i=1}^{m_u} z_i^{-\theta} \log(z_i) e^{-\beta z_i^{-\theta}} \left(1 - \beta z_i^{-\theta} \right) + \sum_{i=1}^{m_u} \varepsilon_i \left(e^{-\beta z_i^{-\theta}} - \right. \\
 & \left. 1 \right) z_i^{-\theta} \log(z_i) e^{-\beta z_i^{-\theta}} \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right) \left[1 - \beta z_i^{-\theta} \left(1 + \frac{1}{\alpha} e^{-\beta z_i^{-\theta}} z_i^{-\theta} \left(e^{-\beta z_i^{-\theta}} - 2 \right) \right) \right] + \sum_{i=m_u+1}^m [\tau + \\
 & \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] + \log(\alpha) \sum_{i=m_u+1}^m e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} [\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] \left(1 - \right. \\
 & \left. \beta[\tau + \lambda(z_i - \tau)]^{-\theta} \right) + \sum_{i=m_u+1}^m \varepsilon_i [\tau + \lambda(z_i - \tau)]^{-\theta} \log[\tau + \lambda(z_i - \tau)] e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \times \\
 & \left[\frac{\left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 \right) \left(1 - \beta - \frac{\beta}{\alpha} [\tau + \lambda(z_i - \tau)]^{-\theta} \right) - \beta[\tau + \lambda(z_i - \tau)]^{-\theta} \left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2} \right] + \\
 & \varepsilon_m [\tau + \lambda(z_m - \tau)]^{-\theta} \log[\tau + \lambda(z_m - \tau)] e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \times \\
 & \left[\frac{\left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right) \left(1 - \beta - \frac{\beta}{\alpha} [\tau + \lambda(z_m - \tau)]^{-\theta} \right) - \beta[\tau + \lambda(z_m - \tau)]^{-\theta} \left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2} \right], \tag{A7}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \lambda \partial \theta} = & \beta \sum_{i=m_u+1}^m (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta-1} (1 - \theta \log[\tau + \lambda(z_i - \tau)]) \\
 & - \sum_{i=m_u+1}^m \frac{(z_i - \tau)}{[\tau + \lambda(z_i - \tau)]} \\
 & + \log(\alpha) \sum_{i=m_u+1}^m \left\{ \left(-\beta^2 e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-2\theta-1} \log[\tau + \lambda(z_i - \tau)] \right) \right. \\
 & + (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta} e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \log[\tau + \lambda(z_i - \tau)] \\
 & \left. + (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta-1} e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right\} \\
 & + \sum_{i=m_u+1}^m \frac{\beta \varepsilon_i}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2} \\
 & \times \left(\theta \beta (z_i - \tau) e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} [\tau + \lambda(z_i - \tau)]^{-\theta-1} \log[\tau + \lambda(z_i - \tau)] \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right. \right. \\
 & \left. \left. - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 \right) \right. \\
 & \left. - \theta [\tau + \lambda(z_i - \tau)]^{-\theta-1} (z_i - \tau) e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 \right) \right. \\
 & \left. + \frac{(z_i - \tau) e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 \right)}{[\tau + \lambda(z_i - \tau)]} \right. \\
 & \left. + e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} [\tau + \lambda(z_i - \tau)]^{-\theta-1} \log[\tau + \lambda(z_i - \tau)] \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right. \right. \\
 & \left. \left. - 2 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 3 \right) e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \beta \theta [\tau + \lambda(z_i - \tau)]^{-\theta-1} \right) \\
 & + \frac{\beta \varepsilon_m}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2} \\
 & \times \left(\theta \beta (z_m - \tau) e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} [\tau + \lambda(z_m - \tau)]^{-\theta-1} \log[\tau + \lambda(z_m - \tau)] \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \right. \right. \\
 & \left. \left. - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right) \right. \\
 & \left. - \theta [\tau + \lambda(z_m - \tau)]^{-\theta-1} (z_m - \tau) e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right) \right. \\
 & \left. + \frac{(z_m - \tau) e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right)}{[\tau + \lambda(z_m - \tau)]} \right. \\
 & \left. + e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} [\tau + \lambda(z_m - \tau)]^{-\theta-1} \log[\tau + \lambda(z_m - \tau)] \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \right. \right. \\
 & \left. \left. - 1 \right) \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \right) \left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 3 \right) e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \beta \theta [\tau + \lambda(z_m - \tau)]^{-\theta-1} \right),
 \end{aligned} \tag{A8}$$

$$\begin{aligned}
 \frac{\partial^2 \ell}{\partial \beta^2} = & -\frac{(m_u+m)}{\beta^2} - \log(\alpha) \sum_{i=1}^{m_u} z_i^{-2\theta} e^{-\beta z_i^{-\theta}} - \sum_{i=1}^{m_u} \varepsilon_i z_i^{-2\theta} e^{-\beta z_i^{-\theta}} \left(\alpha e^{-\beta z_i^{-\theta}} - 1 \right) \\
 & + \log(\alpha) \sum_{i=m_u+1}^m [\tau + \lambda(z_i - \tau)]^{-2\theta} e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \\
 & + \sum_{i=m_u+1}^m \varepsilon_i \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 [\tau + \lambda(z_i - \tau)]^{-\theta} \frac{\left(\left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2 + e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2} \\
 & + \frac{\varepsilon_m \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) [\tau + \lambda(z_m - \tau)]^{-\theta} \left(\left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2 + e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2},
 \end{aligned} \tag{A9}$$

$$\begin{aligned} \frac{\partial^2 \ell}{\partial \beta \partial \lambda} = & -\theta \sum_{i=m_u+1}^m (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta-1} \\ & -\theta \log(\alpha) \sum_{i=m_u+1}^m \left((z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta-1} e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right. \\ & \left. + \beta [\tau + \lambda(z_i - \tau)]^{-2\theta-1} (z_i - \tau) e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} \right) \\ & + \sum_{i=m_u+1}^m \frac{\beta \varepsilon_i}{\alpha} (z_i - \tau) [\tau + \lambda(z_i - \tau)]^{-\theta-1} \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2 \frac{\left(\alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 + 1 \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2} \\ & + \frac{\varepsilon_m \beta}{\alpha} (z_m - \tau) \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2 [\tau + \lambda(z_m - \tau)]^{-\theta-1} \frac{\left(\alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 + 1 \right)}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2}, \end{aligned} \tag{A10}$$

$$\begin{aligned} \frac{\partial^2 \ell}{\partial \lambda^2} = & \beta(\theta - 1) \sum_{i=m_u+1}^m (z_i - \tau)^2 [\tau + \lambda(z_i - \tau)]^{-\theta-2} - (\theta + 1) \sum_{i=m_u+1}^m \frac{(z_i - \tau)^2}{[\tau + \lambda(z_i - \tau)]^2} + \beta \theta \log(\alpha) \sum_{i=m_u+1}^m (z_i - \tau) \\ & e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} [\tau + \lambda(z_i - \tau)]^{-2(\theta-1)} \left([\tau + \lambda(z_i - \tau)]^{-(\theta-1)} + 1 \right) + \sum_{i=m_u+1}^m \theta \varepsilon_i \beta^2 (z_i - \tau) \\ & \tau^2 e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} [\tau + \lambda(z_i - \tau)]^{-\theta-1} \left(e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right) \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 2 \\ & \left(\frac{1 + [\tau + \lambda(z_i - \tau)]^{-(\theta+1)} + e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} + [\tau + \lambda(z_i - \tau)]^{-1}}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_i - \tau)]^{-\theta}} - 1 \right)^2} \right) + \varepsilon_m \beta^2 (z_m - \tau)^2 [\tau + \lambda(z_m - \tau)]^{-\theta-1} \\ & \left(e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right) \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 2 \left(\frac{1 + [\tau + \lambda(z_m - \tau)]^{-(\theta+1)} + e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} + [\tau + \lambda(z_m - \tau)]^{-1}}{\left(1 - \alpha e^{-\beta[\tau + \lambda(z_m - \tau)]^{-\theta}} - 1 \right)^2} \right). \end{aligned} \tag{A11}$$

References

1. Rahman, A.; Sindhu, T.N.; Lone, S.A.; Kamal, M. Statistical inference for Burr Type X distribution using geometric process in accelerated life testing design for time censored data. *Pak. J. Stat. Oper. Res.* **2020**, *16*, 577–586. [\[CrossRef\]](#)
2. Zhang, X.; Yang, J.; Kong, X. Planning constant-stress accelerated life tests with multiple stresses based on D-optimal design. *Qual. Reliab. Eng. Int.* **2021**, *37*, 60–77. [\[CrossRef\]](#)
3. Dusmez, S.; Akin, B. Remaining useful lifetime estimation for degraded power MOSFETs under cyclic thermal stress. In Proceedings of the 2015 IEEE Energy Conversion Congress and Exposition (ECCE), Montreal, QC, Canada, 20–24 September 2015; pp. 3846–3851. [\[CrossRef\]](#)
4. Stojadinovic, N.; Dankovic, D.; Manic, I.; Davidovic, V.; Djoric-Veljko, S.; Golubovic, S. Impact of Negative Bias Temperature Instabilities on Lifetime in p-channel Power VDMOSFETs. In Proceedings of the 2007 8th International Conference on Telecommunications in Modern Satellite, Cable and Broadcasting Services, Nis, Serbia and Montenegro, 26–28 September 2007; pp. 275–282. [\[CrossRef\]](#)
5. Alotaibi, R.; Mutairi, A.; Almetwally, E.M.; Park, C.; Rezk, H. Optimal Design for a Bivariate Step-Stress Accelerated Life Test with Alpha Power Exponential Distribution Based on Type-I Progressive Censored Samples. *Symmetry* **2022**, *14*, 830. [\[CrossRef\]](#)
6. Hassan, A.S.; Nassr, S.G.; Pramanik, S.; Maiti, S.S. Estimation in constant stress partially accelerated life tests for Weibull distribution based on censored competing risks data. *Ann. Data Sci.* **2020**, *7*, 45–62. [\[CrossRef\]](#)
7. Rabie, A. E-Bayesian estimation for a constant-stress partially accelerated life test based on Burr-X Type-I hybrid censored data. *J. Stat. Manag. Syst.* **2021**, *24*, 1649–1667. [\[CrossRef\]](#)
8. Goel, P.K. Some Estimation Problems in the Study of Tampered Random Variables. Ph.D. Thesis, Department of Statistics, Carnegie Mellon University, Pittsburgh, Pennsylvania, 1971.
9. DeGroot, M.H.; Goel, P.K. Bayesian estimation and optimal designs in partially accelerated life testing. *Nav. Res. Logist.* **1979**, *26*, 223–235. [\[CrossRef\]](#)
10. Rahman, A.; Lone, S.A.; Islam, A. Analysis of exponentiated exponential model under step stress partially accelerated life testing plan using progressive type-II censored data. *Investig. Oper.* **2019**, *39*, 551–559.
11. Epstein, B. Truncated life tests in the exponential case. *Ann. Math. Stat.* **1954**, *25*, 555–564. [\[CrossRef\]](#)

12. Balakrishnan, N.; Kundu, D. Hybrid censoring: Models, inferential results and applications. *Comput. Stat. Data Anal.* **2013**, *57*, 166–209. [[CrossRef](#)]
13. Balakrishnan, N. Progressive censoring methodology: An appraisal. *Test* **2007**, *16*, 211–296. [[CrossRef](#)]
14. Balakrishnan, N.; Cramer, E. *The Art of Progressive Censoring: Applications to Reliability and Quality, Statistics for Industry and Technology*; Springer: New York, NY, USA, 2014.
15. Kundu, D.; Joarder, A. Analysis of Type-II progressively hybrid censored data. *Comput. Stat. Data Anal.* **2006**, *50*, 2509–2528. [[CrossRef](#)]
16. Ng, H.K.T.; Kundu, D.; Chan, P.S. Statistical analysis of exponential lifetimes under an adaptive Type-II progressive censoring scheme. *Nav. Res. Logist. (NRL)* **2009**, *56*, 687–698. [[CrossRef](#)]
17. Lin, C.T.; Ng, H.K.T.; Chan, P.S. Statistical inference of Type-II progressively hybrid censored data with Weibull lifetimes. *Commun. Stat. Meth.* **2009**, *38*, 1710–1729. [[CrossRef](#)]
18. Ismail, A.A. Inference for a step-stress partially accelerated life test model with an adaptive Type-II progressively hybrid censored data from Weibull distribution. *J. Comput. Appl. Math.* **2014**, *260*, 533–542. [[CrossRef](#)]
19. Almetwally, E.M.; Almongy, H.M.; Rastogi, M.K.; Ibrahim, M. Maximum product spacing estimation of Weibull distribution under adaptive type-II progressive censoring schemes. *Ann. Data Sci.* **2020**, *7*, 257–279. [[CrossRef](#)]
20. Hemmati, F.; Khorram, E. Statistical analysis of the lognormal distribution under type-II progressive hybrid censoring schemes. *Commun. Stat. Simulat. Comput.* **2013**, *42*, 52–75. [[CrossRef](#)]
21. Sobhi, M.M.A.; Soliman, A.A. Estimation for the exponentiated Weibull model with adaptive Type-II progressive censored schemes. *Appl. Math. Model.* **2016**, *40*, 1180–1192. [[CrossRef](#)]
22. Zhang, C.; Shi, Y. Estimation of the extended Weibull parameters and acceleration factors in the step-stress accelerated life tests under an adaptive progressively hybrid censoring data. *J. Stat. Comput. Simulat.* **2016**, *86*, 3303–3314. [[CrossRef](#)]
23. Nassr, S.G.; Almetwally, E.M.; El Azm, W.S.A. Statistical inference for the extended Weibull distribution based on adaptive type-II progressive hybrid censored competing risks data. *Thail. Stat.* **2021**, *19*, 547–564.
24. Nassar, M.; Nassr, S.G.; Dey, S. Analysis of burr Type-XII distribution under step stress partially accelerated life tests with Type-I and adaptive Type-II progressively hybrid censoring schemes. *Ann. Data Sci.* **2017**, *4*, 227–248. [[CrossRef](#)]
25. Abo-Kasem, O.E.; Almetwally, E.M.; Abu El Azm, W.S. Inferential Survival Analysis for Inverted NH Distribution Under Adaptive Progressive Hybrid Censoring with Application of Transformer Insulation. *Ann. Data Sci.* **2022**, 1–48. [[CrossRef](#)]
26. Alam, I.; Ahmed, A. Parametric and Interval Estimation Under Step-Stress Partially Accelerated Life Tests Using Adaptive Type-II Progressive Hybrid Censoring. *Ann. Data Sci.* **2020**, 1–13. [[CrossRef](#)]
27. Almongy, H.M.; Almetwally, E.M.; Alharbi, R.; Alnagar, D.; Hafez, E.H.; Mohie El-Din, M.M. The Weibull generalized exponential distribution with censored sample: Estimation and application on real data. *Complexity* **2021**, *2021*, 6653534. [[CrossRef](#)]
28. Selim, M.A. Estimation and prediction for Nadarajah-Haghighi distribution based on record values. *Pak. J. Stat.* **2018**, *34*, 77–90. [[CrossRef](#)]
29. Haj Ahmad, H.; Salah, M.M.; Eliwa, M.S.; Ali Alhussain, Z.; Almetwally, E.M.; Ahmed, E.A. Bayesian and non-Bayesian inference under adaptive type-II progressive censored sample with exponentiated power Lindley distribution. *J. Appl. Stat.* **2022**, *49*, 2981–3001. [[CrossRef](#)] [[PubMed](#)]
30. Basheer, A.M. Marshall-Olkin alpha power inverse exponential distribution: Properties and applications. *Ann. Data Sci.* **2019**, *9*, 301–313. [[CrossRef](#)]
31. Neyman, J. Outline of a theory of statistical estimation based on the classical theory of probability. *Philos. Trans. R. Soc. Lond.-Ser. A Math. Phys. Sci.* **1937**, *236*, 333–380.
32. Efron, B. Bootstrap Methods: Another Look at the Jackknife. *Ann. Stat.* **1979**, *7*, 1–26. [[CrossRef](#)]
33. Gelman, A.; Carlin, J.B.; Stern, H.S.; Rubin, D.B. *Bayesian Data Analysis*, 2nd ed.; Chapman and Hall/CRC: Boca Raton, FL, USA, 2004. [[CrossRef](#)]
34. Dey, S. Bayesian estimation of the shape parameter of the generalized exponential distribution under different loss functions. *Pak. J. Stat. Oper. Res.* **2010**, *6*, 163–174. [[CrossRef](#)]
35. Burkschat, M.; Cramer, E.; Kamps, U. On optimal schemes in progressive censoring. *Stat. Probab. Lett.* **2006**, *76*, 1032–1036. [[CrossRef](#)]
36. Burkschat, M.; Cramer, E.; Kamps, U. Optimality criteria and optimal schemes in progressive censoring. *Commun. Stat.—Theory Methods* **2007**, *36*, 1419–1431. [[CrossRef](#)]
37. Burkschat, M. On optimality of extremal schemes in progressive type II censoring. *J. Stat. Plan. Inference* **2008**, *138*, 1647–1659. [[CrossRef](#)]
38. Pradhan, B.; Kundu, D. On progressively censored generalized exponential distribution. *TEST* **2009**, *18*, 497–515. [[CrossRef](#)]
39. Elshahhat, A.; Rastogi, M.K. Estimation of parameters of life for an inverted Nadarajah-Haghighi distribution from type-II progressively censored samples. *J. Indian Soc. Probab. Stat.* **2021**, *22*, 113–154. [[CrossRef](#)]
40. Long, C.; Chen, W.; Yang, R. Ratio estimation of the population mean using auxiliary information under the optimal sampling design. *Probab. Eng. Inf. Sci.* **2022**, *36*, 449–460. [[CrossRef](#)]
41. Badar, M.G.; Priest, A.M. Statistical aspects of fibre and bundle strength in hybrid composites. In *Progress in Science and Engineering Composites*; Hayashi, T., Kawata, K., Umekawa, S., Eds.; ICCM-IV: Tokyo, Japan, 1982; pp. 1129–1136.
42. Chen, G.; Balakrishnan, N. A general-purpose approximate goodness-of-fit test. *J. Qual. Technol.* **1995**, *27*, 154–161. [[CrossRef](#)]

43. Pakyari, R.; Balakrishnan, N. A general-purpose approximate goodness-of-fit test for progressively type-II censored data. *IEEE Trans. Reliab.* **2012**, *61*, 238–244. [[CrossRef](#)]
44. El-Din, M.M.; Abu-Youssef, S.E.; Ali, N.S.; Abd El-Raheem, A.M. Estimation in constant-stress accelerated life tests for extension of the exponential distribution under progressive censoring. *Metron* **2016**, *74*, 253–273. [[CrossRef](#)]
45. Abd El-Raheem, A.M.; Almetwally, E.M.; Mohamed, M.S.; Hafez, E.H. Accelerated life tests for modified Kies exponential lifetime distribution: Binomial removal, transformers turn insulation application and numerical results. *AIMS Math.* **2021**, *6*, 5222–5255. [[CrossRef](#)]
46. Dimitrova, D.S.; Kaishev, V.K.; Tan, S. Computing the Kolmogorov-Smirnov Distribution When the Underlying CDF is Purely Discrete, Mixed, or Continuous. *J. Sta. Softw.* **2020**, *95*, 1–42. [[CrossRef](#)]